

A STUDY OF FAR INFRARED LOOP AT -61° LATITUDE

University of South Carolina
Columbia

Abhinna Rajbanshi
February, 2021

CONTENT

- Background
- Objective
- Theory
- Methodology
- Result & Discussion
- Conclusion
- Reference

BACKGROUND

1. Interstellar Medium

- ❖ Matter and radiation exist in the space between stars.
- ❖ Gases
- ❖ Dusts
- ❖ Cosmic rays
- ❖ Magnetic field

BACKGROUND

2. Infrared Astronomy

- ❖ Electromagnetic Radiation (gamma rays, X-rays, ultraviolet, visible, infrared, microwave and radio wave)
- ❖ Detection and Study of Infrared radiation.
- ❖ Every object that has temperature radiates in infrared.

| Spectral Region | Wavelength Range (microns) | Temperature Range (Kelvin) | What We See |
|-----------------|----------------------------|----------------------------|---|
| Near- Infrared | (0.7-1) to 5 | 740 to (3000-5200) | Color red star, Red Giants Dust is transparent |
| Mid- Infrared | 5 to (25-40) | (92.5-140) to 740 | Planets, comets and asteroid Dust warmed by starlight |
| Far- Infrared | (25-40) to(200-350) | (10.6-18.5) to (92.5-140) | Emission from cold dust Central region of galaxies Very cold molecular clouds |

Source: NASA IPAC “cool cosmos” site

BACKGROUND

3. IRAS Survey

- ❖ The first ever telescope to perform a survey of entire night sky at infrared wavelength.
- ❖ A joint project of the United States (NASA), the Netherlands (NIVR) and United Kingdom (SERC)



fig; Artist's Depiction of IRAS ,

OBJECTIVES

- ❖ **To find an isolated cavity and its possible association using IRAS surveys**
- ❖ **To study the physical properties (dust mass, dust temperature and Planck's function etc.) of cavity**
- ❖ **The stability and the evolution of the cavity will be discussed using published literatures.**

THEORY

1. Far Infrared-loop

- ❖ Cosmic bubble bath (arc, loops, filament, cavity, shells)
- ❖ Supernova Explosion & Stellar winds of associations
- ❖ Supersonic Turbulence may play important role in ruling of structure of ISM, creating clouds and cavities.
- ❖ Kiss et al. (2004) catalogued 145 loops in 2nd galactic Quadrant.
- ❖ Koenyves et al. (2007) extended the search in 1st, 3rd and 4th Quadrant, found 317 loops.
- ❖ These two works provide the catalogued of Far-Infrared loops in the Galaxy.
- ❖ Also known as KK loops.

THEORY

4. Dust Color Temperature

- ❖ It is generally equivalent temperature of dust due to its color (wavelength)
- ❖ According to Schnee et al. (2005), the flux density at λ_i is given by

$$F_i = \left[\frac{2hc}{\lambda_i^3 \left(e^{\frac{hc}{\lambda_i k T_d}} - 1 \right)} \right] N_d \alpha \lambda_i^{-\beta} \Omega_i$$

where α is a constant that relates the flux to optical depth of the dust, Ω_i is the solid angle subtended by the detector

- ❖ According to Wood et al. (2005), $\Omega(60) = \Omega(100)$ and dusts are in thermal equilibrium, we have the formulas for T_d as

$$T_d = - \frac{96}{\ln\{R \times 0.6^{(3+\beta)}\}} \quad \text{Where, } R = \frac{F(60\mu\text{m})}{F(100\mu\text{m})}$$

Here β is the emissivity spectral index depends on dust, grain properties.

THEORY

5. Dust Mass

- ❖ The flux density from a cloud at distance D containing N spherical dust grain of size a , cross section σ and emissivity $Q(\nu)$ is given as

$$F(\nu) = N \left[\frac{\sigma}{D^2} \right] Q(\nu) B(\nu, T)$$

Suppose v be the volume of each dust grain, then total volume of the cloud is $V=Nv$, where $v = \frac{4}{3}\pi a^3$, and $\sigma = \pi a^2$.


- ❖ The dust mass is given by

$$M = V\rho = \left[\frac{F(\nu)D^2}{B(\nu, T)} \right] \left[\frac{4a}{3Q(\nu)} \right] \rho$$


- ❖ Young et al. (1993) suggested that $a=0.1 \mu\text{m}$, $\rho=3000\text{kgm}^{-3}$ and $Q(\nu)=0.001$ and 0.0046 for $100 \mu\text{m}$ and $60 \mu\text{m}$, then

$$M_{dust} = 0.4 \left[\frac{F_{\nu} D^2}{B(\nu, T)} \right] \quad \text{where,} \quad B(\nu, T) = \frac{2h\nu^3}{c^2} \left[\frac{1}{e^{\frac{h\nu}{kT}} - 1} \right]$$

METHODOLOGY



Coordinates or Source:



RA, Dec

(e.g. "Eta Carinae", "10 45 3.6, -59 41 4.2", or "161.265, -59.685" [omit the quotes])

Surveys: Select at least one survey

SkyView Surveys

Clear Survey Selections

Gamma Ray:

- Fermi 5
- Fermi 4
- Fermi 3
- Fermi 2
- Fermi 1
- EGRET (3D)
- EGRET <100 MeV

Hard X-ray:

- INT GAL 17-35 Flux
- INT GAL 17-60 Flux
- INT GAL 35-80 Flux
- INTEGRAL/SPI GC
- GRANAT/SIGMA
- RXTE Allsky 3-8keV Flux
- RXTE Allsky 3-20keV Flux

X-ray: Swift BAT:

- BAT SNR 14-195
- BAT SNR 14-20
- BAT SNR 20-24
- BAT SNR 24-35
- BAT SNR 35-50
- BAT SNR 50-75
- BAT SNR 75-100

Soft X-ray:

- SwiftXRTCnt
- SwiftXRTEp
- SwiftXRTInt
- HEAO 1 A-2

ROSAT w/sources:

- RASS-Cnt Soft
- RASS-Cnt Hard
- RASS-Cnt Broad
- PSPC 2.0 Deg-Int
- PSPC 1.0 Deg-Int
- PSPC 0.6 Deg-Int
- HRI

ROSAT Diffuse:

- RASS Background 1
- RASS Background 2
- RASS Background 3
- RASS Background 4
- RASS Background 5
- RASS Background 6
- RASS Background 7

UV:

- GALEX Near UV
- GALEX Far UV
- ROSAT WFC F1
- ROSAT WFC F2
- EUVE 83 A
- EUVE 171 A
- EUVE 405 A

Swift UVOT:

- UVOT WHITE Intensity
- UVOT V Intensity
- UVOT B Intensity
- UVOT U Intensity
- UVOT UVW1 Intensity
- UVOT UVM2 Intensity
- UVOT UVW2 Intensity

Optical:DSS:

- DSS
- DSS1 Blue
- DSS1 Red
- DSS2 Red
- DSS2 Blue
- DSS2 IR

Optical:SDSS:

- SDSSg
- SDSSi
- SDSSr
- SDSSu
- SDSSz
- SDSSdr7g
- SDSSdr7i

Other Optical:

- Mellinger Red
- Mellinger Green
- Mellinger Blue
- NEAT
- H-Alpha Comp
- SHASSA H
- SHASSA CC

IR: IRAS:

- IRIS 100
- SFD100m
- SFD Dust Map
- IRAS 12 micron
- IRAS 25 micron
- IRAS 60 micron
- IRAS 100 micron

IR: 2MASS:

- 2MASS-J
- 2MASS-H
- 2MASS-K

IR: UKIDSS:

- UKIDSS-Y
- UKIDSS-I

IR: WISE:

- WISE 3.4
- WISE 4.6

IR: AKARI:

- AKARI N60
- AKARI WIDE S

IR: Planck:

- Planck 857
- Planck 545

IR: WMAP & COBE:


- WMAP ILC
- WMAP Ka

Activate Windows
Go to Settings to activate Windows.

METHODOLOGY

2. Region of Interest

- ❖ Out of 50 IR loops, 4 IR loops are considered for further study (depending upon symmetric and isolated cavity)



METHODOLOGY

- ❖ From these 4 IR loops, we have selected the loop G299+02 as region of interest based on following reasons:
 - Distance
 - Not vigorously studied (known from SIMBAD)
 - Appear to be symmetric and isolated





Fig.: IRAS images of the region of interest at 60 and 100 μm band, where we can see the cavities clearly. The dark region represents the low flux density region.

METHODOLOGY

3. Contour Map



- ❖ Drawn isocontour at different levels so that we can separate regions of maximum and minimum flux density.
- ❖ Contours are drawn as to include maximum features of the region with the help of ALADIN2.5.



METHODOLOGY

4. Study of Flux Density

- ❖ Each pixels of the cavity at 60 and 100 μm are tagged and corresponding flux densities are noted.



5. Calculation

- ❖ Dust Color Temperature

$$T_d = -\frac{96}{\ln\{R \times 0.6^{(3+\beta)}\}}$$

- ❖ Dust Mass

$$M_{dust} = 0.4 \left[\frac{S_\nu D^2}{B(\nu, T)} \right]$$

- ❖ Planck's Function

$$B(\nu, T) = \frac{2h\nu^3}{c^2} \left[\frac{1}{e^{\frac{h\nu}{kT}} - 1} \right]$$

- ❖ Major and Minor diameter

6. Graph & Plots

- ❖ F(60) VS F(100) plot
- ❖ Contour Maps of Dust Color Temperature & Dust Mass
- ❖ Gaussian Plots of Dust Color Temperature & Dust Mass
- ❖ Distribution of Planck's Function along Major & Minor diameter

RESULT & DISCUSS

1. F(60) VS F(100) Plot

- Positive correlation can be seen.
- Lower values of flux density are found in limited region, whereas data are populated in higher values of flux density.
- More scatter from fitted line and more data's seen in mid values of flux density.
- Temperature found to be 23.48 K





Fig.: F(60) vs F(100) plot

RESULT & DISCUSS

3. Contour Maps of Dust Color Temperature & Dust Mass



- Following the trend that higher the dust mass, lower is the dust temperature.



Slide 19

RA1

RAJBANSHI, ABHINNA, 2/4/2021

RESULT & DISCUSS



4. Gaussian Plot of Dust Color Temperature & Dust Mass



- The distribution of dust mass and dust temperature show a deviation from the normal gaussian distribution.

RESULT & DISCUSS

5. Distribution of Planck's Function along Extension & Compression



- The value of Planck's function is dependent on the distance along major and minor diameter of the cavity.

CONCLUSION

- ❖ The dust color temperature is found to be in the range 23.34 ± 0.07 or 24.38 ± 0.45 K. An offset of less than 5 K suggests that the cavity is relatively stable.
- ❖ The distribution of dust color and dust mass showed a deviation from Gaussian distribution, suggesting the effect of external cause, possibly wind blower (e.g. AGB wind, stellar supersonic wind, supernova explosion etc.)
- ❖ The dust color temperature and dust mass show an expected trend: lower the temperature, higher the density (or mass). This suggests homogeneous distribution of matter satisfying cosmological principle.
- ❖ The distribution of Planck's function along the extension and compression of the cavity is found to be non uniform, suggesting that the dust and grains are not in LTE with the radiation field.

REFERENCES

1. G. L. Vogt. Space-based astronomy; An educator guide with activities for science, mathematics, and technology education (2001).
2. L. V. Tóth, S. Zahorecz, C. Kiss; Infrared Astronomy (2013)
3. Cs. Kiss, A. Moor, and L. V. Toth, A&A 418, 131–141(2004)
4. V. Könyves, Cs. Kiss, A. Moór, Z. T. Kiss , and L. V. Tóth, A&A 463, 1227–1234(2007)
5. B. Aryal, A. K. Jha, R. Wberger, Research in A & A, submitted 2016
6. Young K., Phillips T. G., & Knapp G. R., 1993, ApJ, 409, 2, 725
7. <http://www.ipac.caltech.edu/outreach/Edu/importance.html>

Thank you